

Proteinase K

[WBC Home](#)[Manual/Protocol Index](#)[Catalog Index](#)**Manual Page** | [Catalog Page](#)**I.U.B.: 3.4.21.14*****Tritirachium* alkaline proteinase**

Proteinase K (PROK) is a serine endopeptidase with a broad spectrum of action, isolated from the filtrate of the fungus *Tritirachium album limber*.

Characteristics of Proteinase K from *Tritirachium album limber*:

Molecular weight: 28,900 daltons.

pH Optimum: 7.5 - 12, using denatured hemoglobin as substrate.

Stability: Although calcium ions do not affect the enzyme activity, they do contribute to its stability when present at a concentration of 1 - 5 μ moles. An interesting characteristic of proteinase K is that it retains its activity in the presence of sodium dodecyl sulphate (SDS) or urea. (0.5 - 1% SDS and 1 - 4 M urea). Raising the temperature of the reaction from 37°C to 50°C - 60°C can increase the activity several folds. A special feature of proteinase K is its ability to digest native proteins, thereby inactivating enzymes such as DNase and RNase without recourse to a denaturation process.

Specificity: In addition to cleavage of peptide bonds, it is able to catalyze peptide amide hydrolysis. Proteinase K is inactivated by diisopropyl fluorophosphate (DFP) or phenyl methane sulphonyl fluoride (PMSF). Chelating agents such as citrate and EDTA have no effect on the enzyme activity.

Application: Proteinase K is very useful in the isolation of highly native, undamaged DNAs or RNAs, since most microbial or mammalian DNases and RNases are rapidly inactivated by the enzyme, particularly in the presence of 0.5 - 1% SDS.

Storage: Store at 2 - 8°C.

Worthington proteinase K is supplied as a highly purified lyophilized powder. It is tested to be free of DNase and RNase.

Assay

Method: Proteinase K hydrolyzes hemoglobin denatured with urea, and liberates Folin Positive amino acids and peptides, which are determined as tyrosine equivalents. 1 unit releases 1 μ mole of Folin positive amino acid in 10 minutes at 37°C, pH 7.5, using denatured hemoglobin as substrate.

Reagents

0.05 N HCl - Dilute 0.82 ml conc. HCl to 200 ml with reagent grade water.

0.5 M NaOH - Dissolve 4.0 gm NaOH in 200 ml reagent grade water.

Buffer-Substrate - Dissolve 2.0 gm hemoglobin in 35 ml reagent grade water, add 36.0 gm urea and 16

ml 0.5 M NaOH. Stir for 30 - 60 minutes at room temperature. Add 0.618 gm boric acid and stir. Adjust the pH to 7.5 with 5 N HCl and q.s. to 100 ml.

Tyrosine standard (2.5 nmol/L) - Dissolve 45.3 mg tyrosine in 100 ml of 0.05 N HCl.

0.3 M Trichloroacetic acid - Dissolve 9.8 gm trichloroacetic acid in 200 ml reagent grade water.

Folin Reagent - Add 10 ml Folin-Ciocalteus Phenol Reagent to 20 ml reagent grade water.

Enzyme

Dissolve 10 mg lyophilized material in 1 ml reagent grade water. Prepare a 1:1000 dilution with water immediately before use.

Procedure

Label clear glass test tubes for blank, standard, and test. Add 2.5 ml buffer-substrate and incubate for 5 minutes at 37°C. Start reaction by adding 0.2 ml tyrosine standard to the standard tube, 0.2 ml of sample to the test, and 0.2 ml of 0.05 N HCl to the blank. Incubate for 10 minutes at 37°C. Stop reaction by the addition of 5.0 ml trichloroacetic acid. Mix, then add 0.2 ml of sample to the blank and standard, and add 0.2 ml of 0.05 N HCl to the test. Mix and let stand for 10 minutes at room temperature, filter and pipette into test tubes 1.0 ml of filtrate, 2.0 ml of 0.5 N NaOH, and 0.6 ml of Folin Reagent. Mix well. Let stand for 15 minutes and read A_{578} nm.

Calculation

$$\text{Units/mg} = \frac{0.5 \mu\text{moles tyrosine}}{0.2 \text{ ml} \times 10 \text{ min}} \times \frac{(\text{A}_{578} \text{ of sample} - \text{A}_{578} \text{ of blank})}{\text{A}_{578} \text{ of standard}} \times \text{Dilution}$$

References

Bernabeu, C., Conde, P., Vasquez, D., and Ballesta, J.: Peptidyl Transferase of Bacterial Ribosome: Resistance to Proteinase K, *Eur J Biochem* 93, 527, 1979

Betzel, C., Pal, G., Struck, M., Jany, K., and Saenger, W.: Active-Site Geometry of Proteinase K: Crystallographic Study of Its Complex with a Dipeptide Chloromethyl Ketone Inhibitor, *FEBS Lett* 197, 105, 1986

Brdicka, D., and Krebs, W.: Localization of Enzymes by Means of Proteases, *Biochim Biophys Acta* 297, 203, 1972

Broemme, D., Peters, K., Fink, S., and Fittkau, S.: Enzyme-Substrate Interactions in the Hydrolysis of Peptide Substrates by Thermolysin, Subtilisin BPN', and Proteinase K, *Arch Biochem Biophys* 244, 439, 1986

Crowe, J., Cooper, H., Smith, M., Sims, M., Parker, D., and Gewert, D.: Improved Cloning Efficiency of Polymerase Chain Reaction (PCR) Products After Proteinase-K Digestion, *Nucleic Acids Res* 19, 184, 1991

Cryer, D., Eccleshall, R., and Marmur, J.: Isolation of Yeast DNA in Methods in, *Cell Biology* 12, 39, 1975

Cryer, D., Eccleshall, R., and Marmur, J.: Isolation of Yeast DNA, *Methods Cell Biol* 12, 39, 1975

Dattagupta, J., Fujiwara, T., Grishin, E., Lindner, K., Manor, P., Pieniazek, N., Saenger, W., and Suck, D.: Crystallization of the Fungal Enzyme Proteinase K and Amino Acid Composition, *J Mol Biol* 97, 267, 1975

Ebeling, W., Hennrich, N., Klockow, M., Metz, H., Orth, H., and Lang, F.: Proteinase K from *Tritirachium album Limber*, *Eur J Biochem* 47, 91, 1974

Fourcroy, P.: Isolation of Undegraded Polysomes from Radish Cotyledons: Use of Proteinase K and Cycloheximide, *Phytochem* 19, 7, 1980

Gajda, A., Rudenskaya, G., and Stepanov, V.: Isolation and Comparative Properties of Serine Proteinases of the Microscopic Fungi *Trichoderma lignorum* and *Trichoderma koningii*, *Biokhimiia* 46, 2064, 1981

Hansen, J.: Isolation of Higher Molecular Weight DNA from *Bacillus cereus* T Using Proteinase K, *Prep Biochem* 4, 473, 1974

Hilz, H., Wiegers, U., and Adamietz, P.: Stimulation of Proteinase K Action by Denaturing Agents; Application to the Isolation of Nucleic Acids and the Degradation of "Masked Proteins", *Eur J Biochem* 56, 103, 1975

Holm, C., Meeks-Wagner, D., Fangman, W., and Botstein, D.: A Rapid, Efficient Method for Isolating DNA from Yeast, *Gene* 42, 169, 1986

Jany, K., and Mayer, B.: Proteinase K from *Tritirachium Album limber*: Molecular Mass and Sequence Around the Active Site Serine Residue, *Biol Chem Hoppe Seyler* 366, 485, 1985

Jany, K., Lederer, G., and Mayer, B.: Amino Acid Sequence of Proteinase K from the Mold *Tritirachium Album limber*. Proteinase K: A Subtilisin-related Enzyme with Disulfide Bonds, *FEBS Lett* 199, 139, 1986

Jehanli, A., and Hough, D.: Pronase and Proteinase K Digestion of Human Immunoglobulin M, *Mol Immunol* 22, 557, 1985

Kasche, V., Zollner, R., Amneus, H., and Naslund, L.: A Two-Step Procedure for Quantitative Isolation of Pure Double Strand DNA From Animal Tissue and Cell Cultures, *Prep Biochem* 11, 233, 1981

Kraus, E., and Femfert, U.: Proteinase K from the Mold *Tritirachium album Limber*. Specificity and Mode of Action, *Z Physiol Chem* 357, 937, 1976

Lizardi, P., and Engelberg, A.: Rapid Isolation of RNA Using Proteinase K and Sodium Perchlorate, *Anal Biochem* 98, 116, 1979

Lough, J., Wrenn, D., Mizioroko, H., and Auer, H.: Differential Sensitivity of Chicken MM-Creatine Kinase to Trypsin and Proteinase K, *Int J Biochem* 17, 309, 1985

McCormick, J., Larson, L., and Maher, V.: Problems in the Extraction of DNA when Utilizing Pancreatic RNAase and Pronase, *Biochim Biophys Acta* 349, 145, 1974

McGodkin, R.: RNA Extraction by the Proteinase K Method, *Methods Mol Biol* 2, 109, 1984

Mendolsohn, S., and Young, D.: Inhibition of Ribonuclease: Efficacy of Sodium Dodecyl Sulfate, Diethyl Pyrocarbonate, Proteinase K and Heparin Using a Sensitive Ribonuclease Assay, *Biochim Biophys Acta* 519, 461, 1978

Morihara, K., and Tsuzuki, H.: Specificity of Proteinase K from *Tritirachium album Limber* for Synthetic Peptides, *Agric Biol Chem* 39, 1489, 1975

Muller, A., Hinrichs, W., Wolf, W., and Saenger, W.: Crystal Structure of Calcium-free Proteinase K at 1.5-Å Resolution, *J Biol Chem* 37, 23108, 1994

Orstan, A., and Gafni, A.: Inhibition of Proteinase-K by Phosphorylated Sugars, *Biochem Int* 25, 657, 1991

Pachler, A., Banerjee, A., Dattagupta, J., Fujiwara, T., Lindner, K., Pal, G., Suck, D., Weber, G., and Saenger, W.: Three-Dimensional Structure of Fungal Proteinase K Reveals Similarity to Bacterial Subtilisin, *EMBO J* 3, 1311, 1984

Pal, G., Betzel, C., Jany, K., and Saenger, W.: Crystallization of the Bifunctional Proteinase/Amylase Inhibitor PKI-3 and of Its Complex with Proteinase K, *FEBS Lett* 197, 111, 1986

Petsch, D., Deckwer, W., and Anspach, F.: Proteinase K Digestion of Proteins Improves Detection of Bacterial Endotoxins by the Limulus Amebocyte Lysate Assay: Application for Endotoxin Removal from Cationic Proteins, *Anal Biochem* 259, 42, 1998

Rauber, N., Jany, K., and Pfeleiderer, G.: RNase A Digestion by Proteinase K, *Z., Naturforsch* 33, Sect.C, 9, 1978

Roelcke, D., and Uhlenbruck, G.: Proteinase K: A New Serological Effective Protease from Fungi, *Z. Med. Mikrobiol. Imm.* 155, 156, 1969

Veerisetty, V., and Sehgal, O.: Proteinase K - Sensitive Factor Essential for the Infectivity of Southern Bean Mosaic Virus Ribonucleic Acid, *Phytopathology* 70, 282, 1980

Vidales, F., Bernabeu, C., and Ballesta, J.: Peptidyl Transfer Ribonucleic Acid Hydrolase Activity of Proteinase K, *Biochem* 18, 4155, 1979

Wiegers, U., and Hilz, H.: A New Method Using Proteinase K to Prevent mRNA Degradation During Isolation from HeLa Cells, *Biochem Biophys Res Commun* 44, 513, 1971

Wiegers, U., and Hilz, H.: Rapid Isolation of Undegraded Polysomal RNA without Phenol, *FEBS Lett* 23, 77, 1972